

Rec'd PCT/PTO 13 JUL 2004



GB03/000086
PCT/GB 2003/000886
10/501184-
INVESTOR IN PEOPLE

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ
REC'D 13 FEB 2003
WIPO
FUT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated

L. Mahoney

31 JAN 2003

BEST AVAILABLE COPY

14 JAN 2002

The
Patent
Office

15JAN02 E687672-3 000016
P01/7700 0.00-0200704.5

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road
Newport
South Wales
NP9 1RH

1. Your reference

JHB/JLH

2. Patent application number

(The Patent Office will fill in this part)

0200704.5

3. Full name, address and postcode of the or of each applicant (underline all surnames)

BRITANNIA PHARMACEUTICALS LIMITED
41-51 Brighton Road
Redhill,
Surrey RH1 6YS

Patents ADP number (if you know it)

5621230001

If the applicant is a corporate body, give the country/state of its incorporation

GB

4. Title of the invention

Use of Phospholipids in Peritoneal Dialysis

5. Name of your agent (if you have one)

BROOKES BATCHELLOR

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

102-108 CLERKENWELL ROAD
LONDON
EC1M 5SA

Patents ADP number (if you know it)

08142291001 ✓

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

-

-

-

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

-

-

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

Yes

a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 13

Claim(s) 2

Abstract -

Drawing(s) 3

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*) -

Request for preliminary examination and search (*Patents Form 9/77*) -

Request for substantive examination (*Patents Form 10/77*) -

Any other documents -
(*please specify*)

11. I/We request the grant of a patent on the basis of this application.

Signature

John H Blake

Date

11 January 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

John H Blake

01892 510600

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) *If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.*
- b) *Write your answers in capital letters using black ink or you may type them.*
- c) *If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- d) *If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- e) *Once you have filled in the form you must remember to sign and date it.*
- f) *For details of the fee and ways to pay please contact the Patent Office.*

USE OF PHOSPHOLIPIDS IN PERITONEAL DIALYSIS

This invention relates to the use of surface active phospholipids (SAPL) to improve the efficiency of ultrafiltration (UF) in patients on continuous ambulatory peritoneal dialysis (CAPD).

In 1985, Grahame et al (Perit. Dial. Bull. 1985; 5:109-111) identified surface-active phospholipids (SAPL) within the peritoneal cavities of patients on continuous ambulatory peritoneal dialysis (CAPD). This followed the earlier discovery of SAPL in the pleural cavity by Hills et al (J. Appl. Physiol. 1982; 53:463-469) and forming an oligolamellar lining which lubricates the pleural mesothelium. A similar lining has since been demonstrated reversibly bound (adsorbed) to peritoneal mesothelium; while the efficacy of adsorbed peritoneal SAPL to act as a boundary lubricant and release agent has been demonstrated by standard physical tests (Chen and Hills; Aust. N. Z. J. Surg. 2000; 70:443-447).

Grahame's discovery led to a somewhat tenuous link being established between any reduction in ultrafiltration (UF) in CAPD patients and an increasing loss of SAPL in their spent dialysate (Di Paolo et al; Perit. Dial. Bull. 1986; 6:44-45). This finding led to a spate of clinical trials in the late 1980s and early 1990s in which peritoneal surfactant was replenished in patients by spiking dialysis fluid with exogenous SAPL. The wide spectrum of outcomes ranged from several totally negative results to others where UF was increased. In the few studies where theory was discussed, the mechanism was generally attributed to a rather nebulous role for SAPL in eliminating a stagnant liquid layer adjacent to the mesothelium (Brebrowicz et al; Perit. Dial. Bull. 1987; 7:6-9) although, in specific studies, such a fluid boundary layer has been dispelled as offering no significant resistance to mass transfer of solutes in PD (Flessner et al; Am. J. Physiol. 1985; 248:F413-424). Subsequently a study by Beavis et al (J. Am. Soc. Nephrol. 1993; 3:1954-1960) held that there is no relationship between dialysate phospholipid levels and the adequacy of UF, and that there was no support for a rationale for intraperitoneal phosphatidyl choline administration in CAPD patients with poor UF.

The present invention starts from the knowledge (Chen and Hills, above) that there is a lining of surface active phospholipid (SAPL) reversibly bound (adsorbed) to normal peritoneal mesothelium which acts as a boundary lubricant and release agent

preserving mechanical integrity of this epithelial surface. The present invention is based on the finding that indigenous peritoneal SAPL is capable of imparting semipermeability to a surface to which it is adsorbed, leading to the conclusion that adsorbed SAPL imparts to peritoneal mesothelium the semi-permeability vital for UF
5 and that any deficiency in SAPL can compromise UF.

The present invention is based on the use of powder compositions of phospholipids and liquid, semi-liquid or pasty compositions of phospholipids dispersed in a physiologically acceptable carrier to promote UF in CAPD patients by administering
10 the compositions directly into the peritoneal cavity or by addition of the compositions to the dialysate used in CAPD.

SAPL powders as described in WO 99/51244 (Britannia) are easily administered into body cavities such as the peritoneum by simple "puffers" or other gas stream delivery
15 devices, and the indicated SAPLs spread rapidly into inaccessible areas. Other suitable compositions are the liquid and paste SAPL compositions disclosed in US Patent 6133249 (Hills).

In one aspect the present invention provides a method of improving the efficiency or
20 reducing deficiency of ultrafiltration in continuous ambulatory peritoneal dialysis which comprises administering a composition comprising at least one SAPL in powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid into the peritoneal cavity before commencing CAPD or between CAPD
25 sessions.

Thus the SAPL may be introduced during surgery to prepare a patient for CAPD; and/or subsequently through the incision for the CAPD catheter, or through the catheter itself, between CAPD sessions when one batch of dialysis fluid has been removed and before a fresh batch is supplied.
30

In another aspect the present invention provides a method of improving the efficiency or reducing deficiency of ultrafiltration in continuous ambulatory peritoneal dialysis which comprises administering a composition comprising at least one SAPL in
35 powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid (other than saline) into the dialysis fluid before commencing a CAPD session.

In this aspect the SAPL composition is mixed with the dialysis fluid and delivered with the dialysis fluid via the catheter provided for the fluid in a CAPD session.

5 In another aspect the present invention provides the use of at least one SAPL in powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid (other than saline) to prepare a medicament for reducing improving the efficiency or reducing deficiency of ultrafiltration in continuous ambulatory peritoneal dialysis.

10 Examples of SAPLs which may be used in this invention include phosphatidylcholine (PC), in particular as diacyl phosphatidylcholines (DAPCs), e.g. dioleoyl phosphatidylcholine (DOPC); distearyl phosphatidylcholine (DSPC) and dipalmitoylphosphatidyl choline (DPPC). A spreading agent may be included which
15 functions to reduce the melting point of a DAPC so that it rapidly spreads as a thin film at normal body temperature. Suitable spreading agents include phosphatidyl glycerols (PG); phosphatidyl ethanolamines (PE); phosphatidyl serines (PS) and phosphatidyl inositols (PI). Another useful spreading agent is cholesteryl palmitate (CP).

20 The above spreading agents, especially PG, are believed to enhance or potentiate the binding of the DAPC, especially the DPPC, to an epithelial surface. However compositions based on DPPC alone may sometimes be as effective as compositions based on DPPC/PG.

25 Also pastes prepared by dispersing coarse SAPL particles, for example around 10 μ m in size, may be more effective than when using fine SAPL particles, such as around 5 μ m in size. More generally, the powdered SAPL may have a particle size in the range of 0.5 to 100 μ m, more suitably of 0.5 to 20 μ m, preferably 0.5 to 10 μ m.

30 Most suitably the dry SAPL composition is prepared from phosphatidylcholine (PC) and phosphatidyl glycerol (PG), but the invention is not limited solely to use of these lipids. Natural endogenous materials contain neutral lipids, fats, inorganic ions etc, all of which are integral to their form and function, and inclusion of these in formulations for use in the invention is not excluded. Preferred SAPL compositions are synthetic
35 dipalmitoyl phosphatidylcholine (DPPC) co-precipitated from a common solvent system with PG in the weight ratio of 6:4 to 8:2, especially about 7:3. The

composition is advantageously administered as a dry powder since it spreads extremely rapidly on water.

5 The phospholipids used in accordance with the invention have acyl substituents on the phosphatidyl groups. As in their natural counterparts, the acyl groups may comprise identical or different, saturated or unsaturated acyl radicals, generally C14-22, especially C16-20, acyl radicals. Thus the phospholipids may comprise, by way of acyl radicals, the saturated radicals palmitoyl C16:0 and stearoyl C18:0 and/or the unsaturated radicals oleoyls C18:1 and C18:2. Diacyl substitution is preferred and
10 the phospholipids used in the compositions in accordance with the invention more particularly comprise two identical saturated acyl radicals, especially dipalmitoyl and distearoyl, or a mixture of phospholipids in which such radicals predominate, in particular mixtures in which dipalmitoyl is the major diacyl component. Thus PC and PG may be used may be used with the same diacylphosphatidyl profile as in PC and
15 PG extracted from human or animal or vegetable sources, but if synthetic sources are used the dipalmitoyl component may predominate, as in the DPPC mentioned above.

As also mentioned above, the SAPL compositions are most preferably protein free, but in some circumstances the presence of proteins and adjuvants, especially naturally
20 occurring materials from plant or animal sources, or synthetically derived, may be tolerated, especially proteins associated with PC and PG *in vivo* in conjunction with a dry powdered formulation for use in this invention. Especially apoprotein B marginally improves SAPL adsorption, and so may be useful if tolerated in SAPL compositions for human use.

25 DPPC can be prepared synthetically by acylation of glycerylphosphorylcholine using the method of Baer & Bachrea - Can. J. of Biochem. Physiol 1959, 37, page 953 and is available commercially from Sigma (London) Ltd. The PG may be prepared from egg phosphatidyl-choline by the methods of Comfurions et al, Biochem. Biophys
30 Acta 1977,488, pages 36 to 42; and Dawson, Biochem J. 1967,102, pages 205 to 210, or from other phosphatidyl cholines, such as soy lecithin.

When co-precipitated with DPPC from a common solvent such as chloroform, PG forms with DPPC a fine powder which spreads rapidly over the surfaces of the
35 airways and lungs. The most preferred composition of the invention contains DPPC and a phosphatidyl glycerol derived from egg phosphatidyl choline, which results in a

mixture of C16, C18 (saturated and unsaturated) and C20 (unsaturated) acyl groups.

The SAPL compositions preferably used in accordance with the present invention are finely-divided, solid powders and are described in detail in our co-pending PCT applications WO 99/27920 and WO 00/30654, the whole contents of which are incorporated by reference. However in summary, our above applications indicate that an important feature of the SAPL compositions that are usable in the present invention is that they are in the form of a powder, that is, it is in solid form. The "dry" surfactant has a high surface activity.

When the SAPL is dispersed or dissolved in a carrier liquid, the carrier liquid is typically one which is substantially non-volatile or only sparingly volatile at body temperature. Suitable carriers include physiologically acceptable glycols, especially propylene glycol, polyethylene glycols and glycerol.

The SAPL may be dispersed in the carrier so as to form liquid, semi-liquid or pasty compositions. Semi-liquid or paste compositions are preferred.

Pastes can be prepared by simply dispersing a SAPL powder in the carrier, or when appropriate dissolving the SAPL(s) in heated carrier and allowing the SAPL(s) to precipitate as a powder on cooling, preferably at a loading that will form a paste. A thick paste of the SAPL and carrier is ideal to apply to open wounds to which it adheres well. It enables a much higher concentration of the SAPL to be applied to the incision site.

Propylene glycol is especially effective as a carrier because at room temperature SAPL may be dispersed in it as a paste, but at body temperature a mobile solution is formed. A paste of 400 mg/ml of DPPC in propylene glycol has given 93% protection against adhesions in surgical tests, as described in the experiments below.

Also polyethylene glycols may be prepared which are waxy solids at room temperature and liquids at body temperature, such as for example PEG 600.

Various dispersions of SAPLs in propylene glycol are described in US Patent 6133249, the entire contents of which are incorporated herein by reference. Similarly the powder compositions of WO 99/51244 may be dispersed in a carrier such as

propylene glycol, and the entire disclosure of WO 99/51244 is also incorporated herein by reference.

In whichever form it is delivered, preferably the SAPL composition has two components. Suitably the first component of the SAPL comprises one or more compounds selected from the group consisting of diacyl phosphatidyl cholines. Examples of suitable diacyl phosphatidyl cholines (DAPCs), are dioleoyl phosphatidyl choline (DOPC); distearyl phosphatidyl choline (DSPC) and dipalmitoyl phosphatidyl choline (DPPC). Most preferably, the first component is DPPC.

The second component may comprise one or more compounds selected from the group consisting of phosphatidyl glycerols (PG); phosphatidyl ethanolamines (PE); phosphatidyl serines (PS); phosphatidyl inositols (PI) and cholesteryl palmitate (CP).

Phosphatidyl glycerol (PG) is a preferred second component. PG is also a preferred second component because of its ability to form with the first component, especially PC and particularly DPPC, a very finely-divided, dry powder dispersion in air.

The composition advantageously comprises a diacyl phosphatidyl choline and a phosphatidyl glycerol. The phosphatidyl glycerol is advantageously a diacyl phosphatidyl glycerol. The acyl groups of the phosphatidyl glycerol, which may be the same or different, are advantageously each fatty acid acyl groups which may have from 14 to 22 carbon atoms. In practice, the phosphatidyl glycerol component may be a mixture of phosphatidyl glycerols containing different acyl groups. The phosphatidyl glycerol is expediently obtained by synthesis from purified lecithin, and the composition of the acyl substituents is then dependent on the source of the lecithin used as the raw material. It is preferred for at least a proportion of the fatty acid acyl groups of the phosphatidyl glycerol to be unsaturated fatty acid residues, for example, mono-or di-unsaturated C18 or C20 fatty acid residues.

Preferred acyl substituents in the phosphatidyl glycerol component are palmitoyl, oleoyl, linoleoyl, linolenoyl and arachidonoyl. The medicament preferably comprises dipalmitoyl phosphatidyl choline and phosphatidyl glycerol, with the phosphatidyl moiety of the phosphatidyl glycerol advantageously being obtainable from the phosphatidyl moiety of egg lecithin.

The compositions are administered preferably in a dry, finely-divided state, using a delivery device such as described in our above co-pending applications, or by directly introducing the aerosolised powder, e.g. by a tube which may be coated to aid transport of SAPL, into the peritoneal cavity.

5

While not wishing to be limited to the following theory it is believed that, when absorbed (reversibly bound) to the peritoneal mesothelium, SAPL provides a semi-permeable membrane by which the desired dialysis is implemented. The predicated deficiency of SAPL which contributes to poor UF leads to a deficiency in this absorbed semi-permeable lining. This situation may be corrected by administering exogenous SAPL, advantageously in a form which displays two properties. First it spreads rapidly over the surface of the incumbent fluid for widespread distribution throughout the peritoneal cavity. Secondly, it then absorbs to the epithelial surface to repair/fortify the semi-permeable barrier comprising similar material.

15

It is highly desirable that the SAPL should not break down quickly at the surgical site in the body. One of the factors which will reduce the life of a lining or coating of SAPL will be the presence of enzymes, such as phospholipase A, capable of digesting DPPC and/or PG. Such enzymes only attack the laevorotatory (L) form, which constitutes the naturally occurring form. Therefore, it may be preferable to use the dextrorotatory (D) form of the SAPL(s) or at least a racemic mixture, which is obtained by synthetic routes.

The compositions may also include preservatives where appropriate, such as fungicides, bactericides and anti-oxidants

25

The present invention is supported by the following experimental work.

INTRODUCTION

30

It has been previously demonstrated that there is a lining of surface active phospholipid (SAPL) reversibly bound (adsorbed) to normal peritoneal mesothelium which acts as a boundary lubricant and release agent preserving mechanical integrity of this epithelial surface. In reviewing clinical trials of the use of SAPL (alias "surfactant") to restore ultrafiltration (UF) in patients on peritoneal dialysis (PD), we

35

have speculated that the SAPL lining might also be imparting the semi-permeability vital for UF.

In evaluating this hypothesis, SAPL harvested from the spent dialysate of 5 patients with normal UF has been deposited on to a porous inert medium and the resulting 7 'membranes' clamped in an Ussing chamber used as an osmometer. In every 'membrane' a clinical concentration of glucose (2.5%) was able to induce a statistically significant osmotic pressure when dialysed against saline. This proves that human peritoneal SAPL has the physical capability to impart semi-permeability 10 when adsorbed to a surface. This could also explain the high permeability of the natural membrane to lipophilic substances in PD.

We have also demonstrated how synthetic SAPL in the form of dipalmitoylphosphatidylcholine (DPPC) and its admixture with phosphatidyl glycerol 15 (pumactant) imparts greater osmotic pressure and does so in proportion to the glucose gradient. Both pumactant and DPPC in various physical forms have been widely used for two decades with complete safety in the treatment of the respiratory distress syndrome in newborns. As a very fine powder, pumactant offers a potential role in restoring UF if applied during the interdialytic interval.

20 The question of formulation of exogenous SAPL in restoring ultrafiltration is discussed as a complex physico-chemical compromise between the higher surface activity of saturated PC and its lower solubility in water.

25 MATERIALS AND METHODS

Principle

The mechanical base for 'the membrane' is a fine-pore filter paper proven to be totally 30 permeable to glucose, urea and physiologically relevant ions. SAPL is then deposited as a thin coating and the resulting membrane clamped between the two compartments of an Ussing chamber to form an osmometer. Any osmotic pressure (ΔP) generated between the compartments is measured as the difference in hydrostatic pressure needed to balance ΔP and stop further osmosis - see Figure 1. The SAPL is derived 35 from spent dialysate from CAPD patients with normal UF and compared with synthetic surfactants envisaged as possible sources of replenishment of indigenous

SAPL where UF is inadequate. The driving force for generating an osmotic pressure is provided by glucose in concentration gradients used clinically to induce and control UF in CAPD.

5 Materials

The synthetic surface-active phospholipid (SAPL) was either dipalmitoyl phosphatidylcholine (DPPC) purchased from Lipoid GmbH (Ludwigshafen, Germany) or pumactant provided by Britannia Pharmaceuticals Ltd (Redhill, UK).

10 Human peritoneal SAPL was extracted from the spent dialysate of patients exhibiting normal UF using the Folch method (J. Biol. Chem. 1957; 226:497-509). All chemical reagents (chloroform, methanol and acetone) were at least AR grade and purchased from AJAX Chemicals (Auburn, NSW, Australia) or BDH Laboratory Supplies (Poole, UK). Saline and Dianeal-2 dialysis fluids with glucose concentrations of
15 1.5%, 2.5% and 4.25% (Baxter Healthcare, Old Toongabbie, NSW, Australia) provided the concentration gradients for generating osmotic pressure. Dialysis fluid with a glucose concentration of 3.4% was made by proportionally mixing two different dialysis fluids (with glucose concentrations of 2.5%; and 4.25%).

20 Methods

SAPL membranes were made by applying equal volumes of SAPL in chloroform solution on to both sides of a filter paper (0.2 μm , white nylon, Millipore Corporation, Bedford, USA). Osmotic pressure was measured by clamping the SAPL membranes
25 between the two compartments of an Ussing chamber (Jim's Instrument Manufacturing, Inc., Iowa, USA). Osmotic pressure was measured as the difference in hydrostatic pressure of the compartments needed to stop further water transmission across the membrane. The total capacity and contact area of chambers are approximately 0.7 ml and 0.44 cm^2 . SAPL (2.36 mg of DPPC, pumactant or human
30 peritoneal SAPL) and 3.78 mg SAPL (DPPC or pumactant) were used for different experiments. Two vertical tubes with inner diameters of 1.2 mm were connected to the side, of each for measuring osmotic pressure. In the experiments, the left compartment was always filled with saline and the right side with test solution (Dianeal-2 dialysis fluids with different glucose concentrations). The device is
35 illustrated in Figure 1. At the beginning of the experiment the fluid heights indicating pressure were set the same on both sides of the membrane. The whole device was kept

at 37°C in a water bath and the fluid heights indicating pressure difference were measured and recorded until there was no further movement of fluid. At the end of each experiment the osmotic pressure was recorded as the difference in heights between the two fluid columns. The mean and S.E.M. were calculated for every group of data and the one-way ANOVA test was used for statistical analysis.

The whole study was divided into five sections:

Section I:

10 Measurement of osmotic pressure produced by dialysing saline against Dianeal-2 dialysis fluids with 2.5% glucose concentrations against DPPC (2.36 mg per preparation) membrane (N=8).

Section II:

15 Measurement of osmotic pressure produced by dialysing saline against Dianeal-2 dialysis fluid with 2.5% glucose concentration against pumactant (2.36 mg per preparation) membrane (N= 8)

Section III:

20 Measurement of osmotic pressure produced by dialysing saline against Dianeal-2 dialysis fluid with 2.5% glucose concentrations using extracted human peritoneal SAPL (2.36 mg per preparation) membrane (N=7).

25 Section IV:

Measurement of osmotic pressure produced by dialysing saline against Dianeal-2 dialysis fluids with different glucose concentrations (1.5%, 2.5%, 3.4% and 4.25%) against DPPC (3.78 mg per preparation) membrane (N=8).

30

Section V:

Measurement of osmotic pressure produced by dialysing saline against Dianeal-2 dialysis fluid with different glucose concentrations (1-5%, 2.5%, 3.4% and 4.25%) using pumactant (3.78 mg per preparation)

35 membrane (N=8).

RESULTS

In all experiments an osmotic pressure was generated by dialysing any hypertonic dialysate against saline. The results from Sections I, II and III are given in Figure 2 while those from Sections IV and V are compared in Figure 3. The features of these results can be listed as follows:

1. In 7 runs, each using the pooled SAPL harvested from 5 exchanges, an osmotic pressure was always generated by Dianeal-2.

2. Synthetic SAPL was more effective than indigenous peritoneal SAPL with pumactant more effective than DPPC at the same (2.36 mg) thickness - see Figure 2.

3. Thicker membranes (3.78 mg DPPC) were more effective than thinner membranes (2.36 mg DPPC), - see Figure 2.

4. For the same membrane thickness and composition, the osmotic pressure increased with the glucose driving force for osmosis - see Figure 3 - as predicted by the van't Hoff equation governing osmosis.

5. Pumactant was more effective than pure DPPC at each glucose concentration. At glucose concentrations of 2.5% and 3.4%, pumactant membranes generated statistically significant higher osmotic pressures than DPPC membranes ($p < 0.05$).

DISCUSSION

Although the results of this study show convincingly that human peritoneal SAPL imparts semi-permeability to an inert porous base, it does not prove conclusively that it necessarily does the same to peritoneal mesothelium *in vivo*.

However, there are many factors which support this hypothesis. Firstly we have previously demonstrated by epifluorescence microscopy that there is a lining of SAPL adsorbed to parietal peritoneum which is probably oligolamellar in nature, resembling similar linings adsorbed to pleural mesothelium. Secondly, oligolamellar layers of SAPL in the form of liposomes have long been known to be semi-permeable to such low-molecular weight solutes as NaCl. Thirdly, there is the evidence from clinical trials that there is a association between reduction of UF in PD and loss of SAPL in

dialysate. It could be argued that the quantity recovered from spent dialysate does not necessarily reflect the amount of SAPL adsorbed to parietal mesothelium which surface accounts for 85% of dialysis. However we have demonstrated that exogenous SAPL in the form of radiolabelled DPPC does indeed adsorb to parietal mesothelium.

5 This raises the issue of whether the administration of exogenous SAPL should be employed for the restoration of UF in patients who have lost that capability and what insight the adsorption theory may offer in the formulation of exogenous SAPL for this purpose. In addition, an SAPL barrier would help to explain why the peritoneal membrane is an order of magnitude more permeable to lipid-soluble substances than
10 to other solutes.

In attempting to review the many clinical trials of SAPL in improving UF, the most frustrating aspect was the lack of physico-chemical information on the widely diverse range of formulations which have been tested. Adsorption is a specialised branch of
15 physical chemistry in which the Langmuir isotherm relates the quantity of a substance adsorbed to its concentration in the adjacent fluid phase. The two parameters most desirable for high adsorption of any substance to a solid surface are high surface activity and high solubility in the adjacent phase - dialysate in the case of PD. Hence we selected DPPC as one of our exogenous surfactants because it is generally
20 regarded as the most surface-active phospholipid. This did indeed display better semi-permeability when used in the Ussing chamber as displayed in Figure 2.

Unfortunately it is highly insoluble in water as demonstrated by a critical micelle concentration as low as 5×10^{-10} Molar (20). In order to circumvent this problem, and largely to improve spreading, DPPC has been used as an intimate mixture with
25 PG (pumactant) in the use of surfactant in treating neonates born with the respiratory distress syndrome. Hence it could be fortuitous that not only is this mixture easier to dispense in aqueous fluids, but it has demonstrated the best results in its ability to impart semi-permeability - see Figure 3. This is encouraging because, when applied as a fine dry powder to the peritoneum, it offered excellent results in preventing surgical
30 adhesions. It would need to be adsorbed strongly to peritoneal mesothelium in order to act as an effective boundary lubricant and release (anti-stick) agent protecting the peritoneum. This raises the possibility of using the interdialytic interval as an opportunity to replenish SAPL, adsorbed to peritoneal mesothelium and hence restore ultrafiltration - whether prescribed as a dry powder (e.g. pumactant) or dispensed in
35 dialysate.

In conclusion, there is good evidence that adsorbed surface active phospholipid is providing the semi-permeability of the mesothelium vital for ultrafiltration, this mechanism offering a new physico-chemical approach to the formulation of SAPL for restoring ultrafiltration as set out in the present invention.

CLAIMS

1. A method of improving the efficiency or reducing deficiency of ultrafiltration in continuous ambulatory peritoneal dialysis which comprises administering a
5 composition comprising at least one SAPL in powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid into the peritoneal cavity before commencing CAPD or between CAPD sessions.
2. A method of improving the efficiency or reducing deficiency of ultrafiltration
10 in continuous ambulatory peritoneal dialysis which comprises administering a composition comprising at least one SAPL in powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid (other than saline) into the dialysis fluid before commencing a CAPD session.
- 15 3. Use of at least one SAPL in powder form or dispersed or dissolved in a physiologically acceptable non-volatile carrier liquid (other than saline) to prepare a medicament for reducing improving the efficiency or reducing deficiency of ultrafiltration in continuous ambulatory peritoneal dialysis.
- 20 4. Use or method according to claim 1, 2 or 3 in the SAPL is selected from diacyl phosphatidylcholines (DAPCs), such as dioleoyl phosphatidylcholine (DOPC); distearyl phosphatidylcholine (DSPC) and dipalmitoyl phosphatidylcholine (DPPC).
5. Use or method according to claim 4 in which the SAPL composition further
25 includes a spreading agent such as a phosphatidyl glycerol (PG), phosphatidyl ethanolamine (PE), phosphatidyl serine (PS), phosphatidyl inositol (PI) or cholesteryl palmitate (CP).
6. Use or method according to claim 1, 2 or 3 in which the SAPL composition is
30 a mixture of phosphatidylcholine (PC) and phosphatidyl glycerol (PG).
7. Use or method according to claim 6 in which the SAPL composition is a mixture of dipalmitoyl phosphatidylcholine (DPPC), or a phosphatidylcholine blend (PC) which is predominantly dipalmitoyl phosphatidylcholine (DPPC). and
35 phosphatidyl glycerol (PG)

8. Use or method according to any preceding claim in which the carrier is glycerol, propylene glycol, or a polyethylene glycol.

9. Use or method according to claim 8 in which the carrier is propylene glycol.

5

10. Use or method according to any preceding claim in which the SAPL/carrier is in the form of a paste

1/3




Figure 1

213




Figure 2.

3/3




Figure 3

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.